



RESEARCH ARTICLE

Open Access

Phytochemical Screening of *Azadirachta Indica* (Neem) Leaves: Revealing Health and Environmental Benefits

Olayiwola Bolaji¹, Yisa Adeniyi Abolade², Sarah Aduwa³, Amarachukwu Bernaldine Isiaka⁴, Olamide Durodola⁵, Abidemi Adeoye⁶, Tola Ojemai Adeoye⁷, Tasha Siame^{8*} and Muawiya Baba Aminu⁹

¹Department of Medicine, New Jersey Medical School, Rutgers University, Newark, USA

²Georgia State University, Atlanta Georgia, USA

³University of west Florida, Earth and Environmental Sciences, USA

⁴Qatar Airways Medical Division, USA

⁵Auburn University, Alabama, USA

⁶Indiana University Bloomington, O'Neill School of Public and Environmental Affairs, USA

⁷International American University, Los Angeles, California, USA

⁸Faculty of Public and Environmental Affairs, Indiana University Bloomington, IN, USA

⁹Department of Geology, Faculty of Sciences, Federal University Lokoja, Kogi State, Nigeria

ABSTRACT

The phytochemical screening of (*Azadirachta indica*) Neem leaf extract was conducted to identify and quantify the presence of bioactive compounds. The study conducted various tests to assess the phytochemical components. About 1kg of matured fresh leaves of (*Azadirachta indica*) neem was collected from five (5) different location in Bauchi metropolis, these locations are; Kobi Street, Gida Dubu, Yelwan Tudu, Wuntin dada and Federal Low-cost areas in Bauchi metropolis, Bauchi State, Nigeria. The leaves of the plant, which were ground into a fine powder using a mortar and pestle. An ethylacetate solvent was used to derive the *A. indica* leaf extract. The results of the research shows that the presence of alkaloids, saponins, phenols, and cardiac glycosides in the neem leaf extract is significant, and flavonoids, terpenoids, and steroids were not detected in the neem leaf extract. The presence of alkaloids, saponins, phenols, and cardiac glycosides suggests that neem leaves may hold promise in traditional and contemporary medicinal practices. Alkaloids, known for their pharmacological significance, may contribute to the plant's therapeutic properties. Saponins, with their frothing and foaming characteristics, have been associated with immunomodulatory effects and their potential in agrochemical applications. Phenolic compounds, as evidenced by the positive response to the Ferric chloride test, indicate the antioxidant potential of neem leaves, which can combat oxidative stress. Conversely, the absence of flavonoids, terpenoids, and steroids suggests that these specific phytochemical classes may be limited in neem leaves. These findings open avenues for further research and utilization of neem as a valuable natural resource.

ARTICLE HISTORY

Received January 01, 2024

Accepted January 09, 2024

Published March 31, 2025

KEYWORDS

Azadirachta Indica, Mortar and Pestle, Ethylacetate Solvent, Cardiac Glycosides

Introduction

Medicinal plants are generated from a variety of plant groups and families and are used as essential oils, plant extract, or both [1]. Many bioactive chemicals with antibacterial and antifungal properties may be found in plant extracts from stem plants, rhizomes, bark, stem, and root [2]. According to Khanal the biological capabilities of medicinal plants are attributed to numerous active components. The primary aim of this research

is to employ an ethyl acetate extract to detect the phytochemical features of the (*Azadirachta indica*) Neem leaves in the study area. The specific

objective of the study was; To identify and quantify the phytochemicals present in neem leaves, to assess the potential health and environmental benefits of the detected phytochemicals.

Contact Tasha Siame, Faculty of Public and Environmental Affairs, Indiana University Bloomington, IN, USA.

© 2025 The Authors. This is an open access article under the terms of the Creative Commons Attribution NonCommercial ShareAlike 4.0 (<https://creativecommons.org/licenses/by-nc-sa/4.0/>).

Neem (*Azadirachta indica*) is a tree in the Meliaceae family. Neem is known by the Sanskrit name "arista," which means "perfect, complete, and perishable" [3]. It is an attractive perennial tree that is native to the Indian region. In addition, it is grown in numerous Latin American nations, Southeast Asia, Australia, East and Sub-Saharan Africa, Fiji, and Mauritius [4]. In the mid-1900s, a large amount of chemical research was done on the byproducts of the neem tree. The United Nations decided to name the amazing neem tree as the Tree of the 21st Century [5]. The most often prescribed remedies in historical medical books for digestive disorders, diarrhea and intestinal infections, skin ulcers, and malaria are extracts from the Neem tree (*A. indica*), also known as "Dogonyaro" in Nigeria [6]. The most often prescribed remedies in historical medical books for illnesses such as diarrhea and intestinal infections, skin ulcers, and malaria are extracts from the Neem tree (*A. indica*), also known as "Dogonyaro" in Nigeria [6].

Global, agricultural, environmental, and health issues have long been addressed using neem [7,8]. Its leaves can be taken as medication to treat eczema, diabetes, and lower fever. Neem leaves are used to produce toothbrushes, and the plant's roots are insect- and disease-repelling. Neem tree seeds are extremely rich in oil. Many illnesses, including diabetes and TB, are treated with neem oil, which is also used as a lubricant, pesticide, and medication [6,9]. Neem leaf extracts have been used in Africa to create a variety of therapeutic medicines. Additionally, it has been demonstrated that various neem parts, including the leaf, bark, and seed oils, have a wide range of pharmacological activities, such as anti-inflammatory, anti-mutagenic, anti-carcinogenic, antioxidant, antihyperglycemic, antiulcer, and anti-diabetic features [10].

The robust, evergreen neem (*Azadirachta indica*) tree can be found mainly in tropical and subtropical regions of the world, primarily in Africa and Asia. It possesses thick, round, pinnate leaves and bark that ranges from brown to dark gray. They range in size from medium to big [11]. The reason for their survival in dry and semi-arid regions of the planet is their vast deep root system. Plants generate molecules known as phytochemicals that help protect against bacteria, viruses, fungus, and weariness in insects and other animals. The name originates from Greek (phyton)

'plant', which is thought to be responsible for protective health benefits [12]. Numerous physiologically active substances, such as alkaloids, flavonoids, triterpenoids, phenolic compounds, carotenoids, steroids, and ketones, may be isolated from neem and added to the chemical contents. Azadirachtin is truly a combination of seven isomeric molecules identified as azadirachtin A-G, and azadirachtin E is more effective [13]. Other molecules that have biological activity are salannin, volatile oils, meliantriol, and nimbin [2].

As antibiotic resistance is a serious problem, the creation of novel compounds derived from plants may help address the need for new antimicrobial agents with enhanced safety and efficacy [1].

Alkaloids, flavonoids, tannins, saponins, polyphenols, and reducing sugars are among the bioactive substances found in neem extracts' leaf and stem bark, according to Bassey [8]. These

chemicals also have antibacterial properties. Further research by Maria and Romilly revealed the presence of volatile oils, triterpenes, alkaloids, flavonoids, tannins, saponins, steroids, resins, bitter cardiac glycosides, reducing sugar, and flavonoids in both fresh and sun-dried neem leaves [14]. These compounds have a significant potential for antibacterial effects.



Figure 1: *Azadirachta Indica* Leaves

This research is justified by the need to elucidate the phytochemical constituents of *Azadirachta indica* (neem leaves) when extracted using ethyl acetate. The results of this research may contribute to the development of new medicines and the ethical use of this valuable plant in a variety of environments by expanding our understanding of the health, nutritional, and safety benefits of neem leaves.

A total of five (5) sample points were selected based on the geographical representative in each direction (see Figure 2). During the sorting of the sample, out of the 5 samples selected and collected, only one important point (Kobi Street) was selected and proceeded with for the research because of its geographical location in the study boundary line and the highest density of the population in the research area.




Figure 2: Map of the Study Area and the Sample Points [15-17].

Neem tree belongs to the family Meliaceae which is widely distributed in tropical and semitropical areas such as Bangladesh, Pakistan, Nepal and India. Additionally, neem has been introduced to various other tropical and subtropical regions worldwide, including Africa, the Caribbean, and parts of the Americas [3]. The tree grows quickly, reaching a height of 20–23 m tall, its trunk is straight, with a diameter of around 4-5 feet. although it can grow taller under favorable conditions [3].

According to Alzohairy the complex, imparipinnate leaves are made up of five to fifteen lance-shaped, serrated-edged leaflets per leaflet [18]. They are arranged alternately on the branch. Its fruits are green drupes which turn golden yellow on ripening in the months of June– August. The fruits are olive-like drupes that are typically oval or round in shape. These fruits are green when young and turn yellow as they ripen. Each fruit contains a single elongated seed with a kernel

inside [19]. The bark of the neem tree is brown to grayish-brown and is characterized by its rough texture. The root has a shallow but widespread root system, making it well-adapted to arid and semi-arid regions [19]. Table I below shows the taxonomic classification of (*Azadirachta indica*) neem.

Table 1: Taxonomic Position of (*Azadirachta Indica*) Neem

Kingdom	Plantae
Order	Rutales
Suborder	Rutinae
Family	Meliceae
Sub-family	Melioidae
Tribe	Melieae
Genus	<i>Azadirachta</i>
Species	<i>Indica</i>

Azadirachta indica L. shows a therapeutic effect in health management because of its abundance of various types of ingredients. Azadirachtin is the most significant active ingredient; the others include quercetin, sodium nimbin, gedunin, salannin, nimbin, nimbidin, and nimbidol. Ascorbic acid, n-hexacosanol, amino acids, nimbin, nimbanene, 6-desacetylnimbinene, nimbandiol, nimbolide, 7-desacetyl-7-benzoylazadiradione, 7-desacetyl-7-benzoylgedunin, 17-

hydroxyazadiradione, and nimbiol are among the ingredients found in leaves [20]. β -sitosterol, Polyphenolic and Quercetin flavonoids, were extracted from neem fresh leaves and were known to have antibacterial and antifungal properties and seeds hold valuable constituents such as azadirachtin and gedunin. Neem is renowned for its diverse array of bioactive compounds, which contribute to its therapeutic potential [20]. Below are the key phytochemical constituents found in neem leaves, seeds, and other parts of the tree.

Alkaloid

Neem leaves and seeds contain various alkaloids, including nimbin, nimbidin, and nimbinene. These alkaloids have been associated with various pharmacological activities, such as anti-inflammatory, analgesic, and antipyretic effects [21].

Flavonoids

Neem leaves are a good source of flavonoids, including quercetin, kaempferol, and rutin. Flavonoids are well-known for their antioxidant properties, which can help combat oxidative stress and reduce the risk of chronic diseases [22].

Saponin

Neem seeds are particularly rich in saponins. Saponins have demonstrated various biological activities, including anti-inflammatory, immunomodulatory, and antitumor effects [23,24].

Phenolic Compounds

Neem leaves contain phenolic compounds, such as gallic acid and catechin. These compounds contribute to the antioxidant properties of neem, which can safeguard the cells from harm caused by free primitives [25].

Terpenoids

Neem is a source of various terpenoids, including limonoids, triterpenes, and essential oils. Limonoids found in neem, such as azadirachtin, have insecticidal properties and are used in organic pest control [25].

Steroids

Neem seeds contain steroids, including beta-sitosterol. Beta-sitosterol has been studied for its potential cholesterol-lowering effects and anti-inflammatory properties [26].

Cardiac Glycosides

Neem leaves have been found to contain cardiac glycosides. Cardiac glycosides have a well-documented role in cardiac health and have been studied for their potential in managing heart conditions [27].

Tannins

Tannins are also present in neem leaves and seeds. They have astringent properties and can form complexes with proteins, which may contribute to their medicinal use [28].

Potential Health and Environmental Benefits of Neem Phytochemicals

Neem (*Azadirachta indica*) has been utilized traditionally for its various phytochemicals in both medicine and agriculture. Additionally, contemporary research has shed light on numerous applications of neem-derived compounds. Below are the traditional and contemporary uses of neem phytochemicals.

Potential Health Application of Neem Phytochemicals

Neem's bioactive compounds, such as nimbidin and nimbin, have been traditionally applied in Ayurvedic medicine for their antibacterial and antifungal properties. Neem-based formulations have been used to alleviate inflammation and pain in various conditions, including arthritis [29].

Antioxidant and Anti-Inflammatory Effects

Neem's phytochemicals, particularly phenolic compounds, exhibit

strong antioxidant properties. Antioxidants neutralize free radicals in the body, protecting cells from oxidative stress and inflammation [24]. These properties suggest a potential role in preventing and managing inflammatory conditions. Neem extracts, owing to their antimicrobial and anti-inflammatory properties, are used for treating skin conditions like acne, eczema, and psoriasis [30].

Cardioprotective Properties

The presence of cardiac glycosides in neem suggests potential cardioprotective effects. Cardiac glycosides have been studied for their positive impact on heart health, including their use in managing cardiac conditions [27]. Further research can explore neem's role in cardiovascular health.

Antimicrobial and Antibacterial Activity

Alkaloids, identified in moderate concentrations in neem, contribute to its antimicrobial properties. Neem has been traditionally used to treat various infections due to its antibacterial and antifungal

effects [19]. This suggests applications in the development of natural antimicrobial agents. Neem twigs have been used traditionally as a natural toothbrush due to their antibacterial properties, helping to maintain oral hygiene [22,31].

Immune System Modulation

Saponins, present at a low concentration, are known for their immunomodulatory properties. These compounds may play a role in enhancing the immune system's response to infections and diseases [23]. Neem's immunomodulatory potential can be explored for therapeutic applications.

Potential Environmental Applications of Neem Phytochemicals Organic Pest Control

Azadirachtin, a prominent neem-derived compound, is widely used in organic pest control due to its insecticidal properties [25]. Neem extracts can serve as eco-friendly alternatives to synthetic pesticides, minimizing environmental impact.

Phytoremediation

Neem trees are planted in contaminated areas to help purify soil and groundwater due to their ability to absorb and neutralize pollutants. Immunomodulatory Properties: Neem-based products may have applications in immunomodulation and supporting the immune system [23]. Ineffective protein utilization in food that has disastrous effects on the ecosystem due to nitrogen pollution [17,32].

Sustainable Agriculture

Neem-based fertilizers, such as neem cake, provide a sustainable alternative to chemical fertilizers. The organic matter in neem cake enhances soil fertility and promotes sustainable agriculture practices [25].

Biopesticides for Crop Protection

Neem extracts, containing a combination of bioactive compounds, can be formulated into biopesticides. These biopesticides offer environmentally friendly alternatives for crop protection, reducing reliance on synthetic chemicals [25].

A study by provides a comprehensive overview of the phytochemical composition of neem leaves, highlighting the presence of alkaloids, flavonoids, and other bioactive compounds [19]. It discusses the traditional uses and medicinal properties of these phytochemicals. conducted a study that explores the traditional use of neem leaves for their antibacterial and antifungal properties [29]. The paper discusses the potential of neem phytochemicals in combating microbial infections. provides an overview of the history and production of azadirachtin, a prominent neem-derived compound used in organic pest control [25]. The study discusses the applications of neem-based pesticides in agriculture. explore the use of neem trees in phytoremediation for environmental cleanup [33]. The paper discusses how neem's phytochemicals can help purify soil and groundwater in contaminated areas. According to a reseach was conducted to assess the 70% alcoholic neem root bark extract (NRE) in diabetes. The findings indicated that the 800 mg/kg dosage of neem root bark extract produced statistically significant outcomes [34].

Further research was conducted to investigate the pharmacological hypoglycemic action of Azadirachta indica in diabetic rats. The research revealed that the rats' glucose levels were significantly lower in the glucose tolerance test using 250 mg/kg of neem extract than in the control group, and that the administration of Azadirachta indica significantly decreased the glucose levels in the diabetic rats by the fifteenth day [35].

Existing research has predominantly focused on specific classes of phytochemicals present in neem leaves, such as alkaloids, flavonoids, and saponins. However, there is a notable gap in our understanding when it comes to a more comprehensive phytochemical analysis. Neem leaves contain a vast array of bioactive compounds, and a holistic examination of these components is necessary to fully elucidate their potential health benefits. A comprehensive analysis would provide a more nuanced view of the synergistic effects between different compounds and their collective impact. This knowledge gap calls for research that explores the full spectrum of neem leaf phytochemicals to better grasp their therapeutic potential [36].

While individual phytochemicals in neem leaves have been studied, the bioavailability of these compounds and their potential synergistic effects when present together is an area that requires

more attention. Understanding how neem phytochemicals are absorbed and utilized by the body is critical for designing effective interventions. Additionally, the potential synergies between these compounds can enhance or alter their therapeutic properties. Research addressing these bioavailability and synergy aspects will shed light on the practical implications of using neem in healthcare [25].

To establish the therapeutic potential of neem leaves and their phytochemical components in humans, particularly for specific health conditions, well-designed clinical trials are required. While preliminary research shows promise, robust clinical studies are necessary to provide evidence-based insights into the efficacy of neem-based interventions. Clinical trials would bridge the gap between laboratory findings and practical healthcare applications, allowing for more informed and evidence-based recommendations for neem usage [25].

The establishment of quality control measures and standards for neem-based products, such as dietary supplements and cosmetics, is an important yet underdeveloped area. Quality control ensures the consistency and safety of products available

to consumers. It is imperative to create guidelines and regulations that guarantee the quality and efficacy of neem-based products, addressing this crucial aspect of neem utilization. For the long-term availability and preservation of Azadirachta indica, research on sustainable cultivation and harvesting practices is essential. Sustainable practices not only support the ecological balance but also ensure a continuous supply of neem leaves. Research in this domain aims to develop environmentally responsible approaches for neem cultivation [33]. Addressing these knowledge gaps is vital for advancing our understanding of neem's phytochemical composition, its potential health applications, and its sustainable utilization. These research areas will contribute to a more comprehensive and evidence-based approach to the use of neem in various fields, from medicine to agriculture.

Materials and Methods




Figure 3: Research Methodology Flow Chat [17,37,38].

Data Collection and Preparation of Plant Sample

Fresh leaves of Azadirachta indica (neem) was collected from Yelwan Tudu, Federal Low cost, Gida Dubu, Wuntin Dada and Kobi Street of Bauchi, Bauchi State. An approximate 1 kg of fresh, matured leaves of the plant was thoroughly cleaned with tap water, after which they were rinsed with distilled water. The leaves were then allowed to air dry for a week at ambient temperature (27 oC). The sample was then manually pounded into a fine powder using a sterile, clean mortar and pestle to enhance surface area, and the powder was weighed, assembled, and packaged into clean cellophane bags and labeled to avoid mix-ups. The sample was stored in a cool, dry location until needed, while fresh leaves were used promptly after being collected from the tree branches.

Extraction of Plant Material

The extraction was prepared using maceration method. Thus, 100g of the powdered plant material was weighted and extracted with 250ml of ethyl acetate for 72 hours. Proper filtration was carried out using filter paper after which the filtrate was boiled for proper extraction. The extract was tested with several reagents to determine the presence and quantity of Tannin, saponins,

alkaloids, flavonoids, terpenoids, steroids, phenolic compound and cardiac glycoside.

Phytochemical screening of Azadirachta indica Extract

To identify the chemical components of the plant's ethyl acetate extracts, routine phytochemical tests were carried out.

Test for Tannins

A small amount of 0.5g of the crude extract was dissolved in distilled water. To this solution 2 ml of 5% ferric chloride solution was added. Formation of blue, green or violet color indicates presence of phenolic compounds [8].

Test for Saponins

The extract was diluted with distilled water and shaken in a graduated cylinder for 15 minutes. The formation of layer of foam indicates the presence of saponins [39].

Test for Alkaloid

The presence of alkaloids in the sample was shown by the appearance of a dark orange or purple hue when 1 ml of each solvent's extract and 1 ml of Wagner's reagent were added and well mixed [8].

Test for Steroid

The extract was treated with chloroform and filtered. The filtrate was added with few drops of concentrated sulphuric acid, shaken and allowed to stand. If the lower layer turns red, steroid is present [8].

Test for Flavanoids

After extracting around 4 g of neem from fresh and dried leaves that had been shaded, 6 ml of a 50% methanol, ethanol, and acetone solution was added. After warming the solutions, 1.5 g of metal magnesium was added to the solution containing neem leaf extract. Then 4-5 drops of concentrated hydrochloric acid were added [8].

Test for Terpenoids

A reddish-brown color at the interface indicates a good result for the presence of terpenoids. The crude extract was mixed with 2 ml of chloroform, and 3 ml of concentrated sulfuric acid was carefully added to produce a layer [8].

Test for Phenol

Distilled water was used to dissolve a small quantity of extract. A 5% ferric chloride solution (two milliliters) was added to this solution. Phenolic chemicals are present when a blue, green, or violet color forms

Test for Cardiac Glycosides

About 2ml of the crude extract was dissolved in 1ml of glacial acetic acid. A few drops of 5% ferric chloride solution and few drops of concentrated sulphuric acid were added. Formation of green blue precipitate indicate the presence of glycosides [8].

Result and Discussion

Result of Phytochemical Screening of the Ethyl Acetate Extracts

Alkaloids, saponins, phenols, and cardiac glycosides were found in the phytochemical screening of the Azadirachta indica leaf extract, as shown in Table II below. Other components analyzed, including flavonoids, terpenoids, and steroids, were not found in the extract.

Table 2: Phytochemical Screening of Azadirachta Indica Leaf Extract

Phytochemical components	Test	Results
Alkaloid	Wagner's test	++
Flavonoid	Ammonia test	-
Saponin	Frothing/foam test	+
Phenol	Ferric chloride (FeCl3) test	+
Terpenoids	Terpenoid test	-
Steroid	Salkowski's test	-
Cardiac glycoside	Keller-Killiani test	++
Tannins	Ferric chloride	+

Key: - = not detected

+ = low concentration

++ = moderate concentration

+++ = high concentration




Figure 4: A Bar Chart Pictorial Representation of The Phytochemical Screening of Azadirachta Indica Leaf Extract

The bar graph represents the phytochemical screening of Azadirachta indica leaf extract. Each bar corresponds to a phytochemical component, and the height of the bar indicates the concentration level detected in the extract:

"Not detected" is represented by the absence of a bar (see figure 4)

"Low concentration" is indicated by a bar with a single plus sign above it (see figure 4). **"Moderate concentration"** is indicated by a bar with two plus signs above it (see figure 4).

"Moderate concentration" is indicated by a bar with two plus signs above it (see figure 4).

Discussion

Phytochemical Screening of Azadirachta Indica Leaf Extract

The phytochemical analysis of Azadirachta indica (neem) leaves revealed the presence and concentration of various bioactive compounds, including alkaloids, saponins, phenols, and cardiac glycosides, which are important for understanding the potential health and environmental applications of neem. As a result, no flavonoids, terpenoids, or steroids were detected.

The Results Indicate the Following Findings

The presence of alkaloids in ethylacetate neem leaves, as determined by the Wagner's test, is significant at a moderate concentration (++). Alkaloids have been shown to have a variety of biological functions, including analgesic and antibacterial characteristics. In their study on Phytochemical analysis of different extract of Azadirachta indica Leaves, detected alkaloids, tannins, phytosterols, and saponins in all extracts of A indica [6]. This finding is also matching with the research of Madaki et al [40]. who discovered alkaloid and other components in their study of Phytochemical and proximate analyses of methanol leaf extract of Neem Azadirachta indica. Alkaloids are primarily found in plant species, where they often serve as secondary metabolites.

The presence of alkaloids also suggests that A. indica extracts may have biological actions, including anti-cholinergic, anti-tumor, anti-hypertensive, cough expectorant, anesthetic, analgesic, muscle relaxant, anti-pyretic, and anti-malarial [11,41]. Recent research has looked into the therapeutic qualities of neem alkaloids, such as their antibacterial, anti-inflammatory, and immunomodulatory effects [42]. Some alkaloids, for example nicotine and caffeine, are well-known for their effects on the human body and are frequently used recreationally or in foods and beverages [42]. Alkaloids like caffeine can increase heart rate and blood pressure. Prolonged excessive consumption of caffeine may contribute to cardiovascular issues, including arrhythmias, high blood pressure, and an increased risk of heart disease. High doses of certain alkaloids can lead to nervous system disturbances, including anxiety, restlessness, tremors, and in severe cases, seizures or hallucinations [43].

The presence of saponins in neem leaves is considerable, as evidenced by a low concentration (+ frothing/foam test). Saponins can be utilized as natural detergents and emulsifiers, among other things. Saponins are also being investigated as immunomodulatory and anticancer medicines [44]. Saponins are polycyclic aglycones connected to one or more sugar side chains. Saponins have a variety of health benefits, including decreased blood cholesterol, cancer prevention, bone health, and immune system activation. Saponins contain antifungal, antibacterial, and antiprotozoal effects, according to Kumar et al [45]. Aqueous leaf extract of A. indica contained alkaloids and saponins with nematocidal activity [46]. Excessive saponin consumption might cause

gastrointestinal problems. Saponins have a foaming property and can form compounds in the digestive tract with lipids and proteins. This can cause nausea, vomiting, diarrhea, and abdominal cramps [47]. It can also irritate the mucous membranes of the mouth, throat, and gastrointestinal system in some situations. This inflammation can be painful and contribute to gastritis or other inflammatory diseases [48]. It also contains antioxidant properties that can help the body neutralize harmful free radicals. This characteristic could contribute to its potential health benefits, such as resistance to oxidative stress-related illnesses [49]. According to study, some saponins present in ginseng and soy may have anticancer properties. They have the ability to inhibit the growth of cancer cells and induce apoptosis (programmed cell death) in these cells. The presence of saponins in neem leaves is considerable, as evidenced by a low concentration (+ frothing/foam test). Saponins can be utilized as natural detergents and emulsifiers, among other things. Saponins are also being investigated as immunomodulatory and anticancer medicines [44].

The presence of phenols, as demonstrated by a positive ferric chloride (FeCl₃) test result, shows that antioxidant effects are possible. Anti-oxidative action has been attributed to phenolic compounds contained in A. indica leaf extracts, which protects cells and tissues from free radicals [45].

Among the most potent natural antioxidants found in plants are phenolic chemicals. By scavenging free radicals and minimizing oxidative damage, they help protect cells from oxidative stress. This feature has been linked to a variety of health advantages [50]. Phenols have anti-inflammatory characteristics that can aid in the reduction of chronic inflammation. Chronic inflammation has been linked to a variety of disorders, including cardiovascular disease and cancer [51]. Some phenolic chemicals, such as resveratrol (found in red grapes and wine), have been studied for their ability to limit cancer cell development and trigger apoptosis, or programmed cell death [50]. Phenolic chemicals have been linked to cardiovascular benefits, such as lowering the risk of heart disease. They can help lower blood pressure, improve blood vessel function, and lower LDL cholesterol oxidation [52]. Excessive consumption of phenolic compound-rich foods, such as certain fruits and vegetables, can cause digestive discomfort, including abdominal pain and diarrhea. This is especially important for people with sensitivities or allergies [53]. While phenolic

chemicals are generally benign, certain highly concentrated phenols, such as hydroquinone, can be poisonous and hazardous if taken in large quantities [53].

The Keller-Killiani test indicated the presence of cardiac glycosides at a moderate concentration (++) . Cardiac glycosides have been studied for their potential utility in heart disease. They have the potential to make an impact on cardiovascular medicine [42]. Traditional medicine has used cardiac glycosides' cardiotoxic effects for millennia. They increase the force and efficiency of cardiac contractions, which makes them effective in illnesses such as congestive heart failure [54]. Cardiac glycosides alter cardiac muscle contractility, which can help enhance heart pumping efficiency [55]. Recent research has explored the potential anticancer properties of certain cardiac glycosides. These compounds have shown promise in inhibiting the growth of cancer cells and inducing apoptosis [56]. Some cardiac glycosides have been investigated for their anti-inflammatory activities, which can have implications for various inflammatory conditions [57]. One of the primary concerns with cardiac glycosides is their narrow therapeutic window, meaning there is a small margin between the therapeutic dose and a toxic dose. Slight overdosage can lead to serious toxicity [58]. Excessive use of cardiac glycosides can cause severe cardiac toxicity, potentially resulting in arrhythmias, heart block, and even death [59]. Nausea, vomiting, stomach pain, and diarrhea are early signs of cardiac glycoside overdose [58]. Cardiac glycosides can also disrupt other physiological systems, including electrolyte imbalances, renal dysfunction, and hyperkalemia (high potassium in the blood) [59].

The presence of tannins, as revealed by the positive ferric chloride test result, implies that neem leaves contain these astringent chemicals. Tannins have been researched for their antiviral, antibacterial, and antioxidant activities, making them appealing for both medicinal and industrial applications [60]. It is well-known for its powerful antioxidant effects. They can help protect cells and tissues from oxidative damage by scavenging free radicals and avoiding lipid peroxidation [49]. It has also been shown to have anti-inflammatory characteristics, making it potentially effective in decreasing chronic inflammation, which plays a critical role in many chronic diseases [50]. Some tannins have antibacterial properties that help to keep bacteria, fungi, and viruses at bay. Cushnie and Lamb state that this can be effective for food preservation and infection prevention [61]. Tannins, notably those contained in red wine and some fruits, have been linked to cardiovascular benefits. They can improve blood vessel function, lower LDL cholesterol oxidation, and assist regulate blood pressure [52]. Excessive consumption of tannin-rich foods or beverages can cause digestive discomfort, including abdominal pain, nausea, vomiting, and constipation [62]. Tannins are astringent and can cause tooth discoloration and erosion. Tannin-containing beverages such as tea and red wine may have a deleterious impact on tooth health [63].

The absence of flavonoids, as shown by the negative ammonia test result, is quite unexpected. Flavonoids are renowned for their antioxidant and anti-inflammatory qualities, and their presence is frequently associated with plant leaves. However, it is important to remember that the absence of flavonoids in the specific sample

studied does not rule out their presence in other portions of the neem plant or under different conditions [64]. The absence of flavonoids in the studied neem ethyl acetate extract could be attributed to changes in textural, chemical, and nutritional soil variables that vary between geographic areas [65,66]. Flavonoids are renowned for their strong antioxidant properties, which help protect cells and tissues from oxidative damage by neutralizing harmful free radicals [50]. Flavonoids have been shown to exhibit anti-inflammatory properties, making them valuable in reducing chronic inflammation, a key factor in many chronic diseases [50]. Some flavonoids have natural antimicrobial properties and can help inhibit the growth of bacteria, fungi, and viruses, which can be beneficial for food preservation and combatting infections [61]. Excessive consumption of flavonoid-rich foods or supplements may cause digestive discomfort such as abdominal pain, nausea, vomiting, and diarrhea [67]. While rare, some people may be allergic to particular flavonoids and have skin rashes and itching after consuming them [68]. Flavonoids, particularly those found in tea and certain fruits, have been demonstrated to restrict the absorption of non-heme iron (the iron found in plant-based meals) in the stomach. This could result in iron insufficiency in persons who don't obtain enough iron from their diet [68].

The absence of terpenoids and steroids, as evidenced by the negative findings in the terpenoid and Salkowski's tests, shows that neem leaves may be deficient in these phytochemical groups. Terpenoids, which are renowned for their varied biological actions, are frequently present in plant essential oils, whereas steroids are key components of many bioactive chemicals [69]. Terpenoids, which include carotenoids and tocopherols, have substantial antioxidant action. They help protect

cells and tissues from oxidative stress by neutralizing damaging free radicals [69]. Certain terpenoids, particularly essential oils containing terpenes, have inherent antibacterial effects. They have the ability to prevent the growth of bacteria, fungus, and other microorganisms [70]. In some cases, the consumption of terpene-rich essential oils or concentrated terpenoid supplements can lead to gastrointestinal discomfort, including nausea and diarrhea [71]. Direct skin contacts with certain terpenes, particularly in essential oils, may cause skin irritation or allergic reactions in sensitive individuals [72].

Plant steroid phytosterols have cholesterol-lowering effects and are frequently added to functional foods. They can lower dietary cholesterol absorption in the intestines [73]. Some plant steroids, such as beta-sitosterol, have been demonstrated to have anti-inflammatory properties. They may help reduce inflammation and are being studied for their possible function in the treatment of illnesses such as arthritis [74]. High dosages of phytosterols or other plant steroids can impair fat-soluble vitamin absorption, particularly vitamins A, D, E, and K [75]. Although it is uncommon, some people are allergic to plant sterols and may develop allergic reactions when eating foods supplemented with high quantities of phytosterols [76].

Conclusion and Recommendations

Conclusion

The phytochemical analysis of Azadirachta indica (neem leaves) revealed the presence of alkaloids, cardiac glycosides, saponins, phenols, and tannins. However, flavonoids, terpenoids, and steroids were either absent or identified in low amounts. These findings reveal that neem leaves contain a variety of bioactive chemicals, including those with potential antibacterial, anti-inflammatory, antiarrhythmic, antiviral, antifungal, cardioprotective, and antioxidant effects, which are consistent with the universal use of neem in herbal medicine. Further study and clinical trials are needed to better understand the precise health consequences and therapeutic applications of the phytochemicals found in neem leaves.

Recommendations

Base on the Findings of the Present Study, the Following Recommendation were Made:

- i. Research efforts should continue to examine and identify the whole spectrum of phytochemicals found in Azadirachta indica (neem leaves) in order to get a comprehensive grasp of the plant's phytochemistry.
- ii. Future research should concentrate on the possible medicinal applications of neem leaves and their phytochemical constituents.
- iii. In light of the moderate concentrations of cardiac glycosides and alkaloids, it is imperative to conduct in-depth toxicological studies.

Clinical trials should be initiated to evaluate the therapeutic potential of neem leaves and their phytochemical components in treating specific health conditions.

References

- [1] Srivastava, Shukla A, Kumar YN. Recent development in plant derived antimicrobial constituents A Review. Journal of Medical and Aromatic Plant Science. 2010; 20: 717- 772.
- [2] Khanal S. Qualitative and quantitative phytochemical screening of Azadirachta indica Juss. Plant Parts. International Journal of Applied Sciences and Biotechnology. 2021; 9: 122-127.
- [3] Girish K, Neem SBS. "A green treasure," Electronic Journal of Biology. 2008; 4: 102-111.
- [4] 4Puvan AA, Irfan M, Rosdan S, Zeehaida M. 'Antifungal effect of Malaysian Neem leaf extract on selected Fungal specie causing Otomycosis in-vitro culture medium', Malaysian Journal of Medicine and Health Sciences. 2015; 11: 69-84.
- [5] Puri H, Neem S. The divine tree; Azadirachta indica. Amsterdam: Harwood Academic Publishers. 1999; 1-3.
- [6] Virshette SJ, Patil MK, Deshmukh AA, Shaikh JR, Dhas MS. Phytochemical constituents of different extracts of Azadirachta indica leaves in urine solvent of a non-pregnant cow. Journal of Pharmacognosy and Phytochemistry. 2020; 9: 1324-1328.
- [7] Santhosh VK, Navartnam V. Neem prehistory to contemporary medicinal uses to humankind. Asian Pacific Journal of Tropical Biomedicine. 2013; 3: 505-519.
- [8] Bassey EE, Mohammed GA, Bala HM, Ogonna US, Yawuri BB, et al. 'Phytochemical analysis and antimicrobial activity of methanolic, ethanolic and acetonic extracts of stem bark and leaf of Neem plant (Azadirachta indica). Journal of Diseases and Medical Plants. 2016; 2: 14-25.
- [9] Kumar AG, Wu CJ, Kumar BJ, Ha DC. Antioxidant activity and quantitative estimation of azadirachtin and nimbin in Azadirachta indica A. Juss grown in foothills of Nepal African Journal of Biotechnology. 2009; 8: 3084-3091.
- [10] Reddy RRY, Krishna KC, Lokanatha O, Mamatha S, Reddy DC. 'Antimicrobial activity of Azadirachta indica (neem) leaf, bark and seed extract', International Journal of Research in Phytochemistry and Pharmacology. 2013; 3: 1-4.
- [11] Susmitha S, Vidyamol KK, Ranganayaki P, Vijayaragavan R. Phytochemical Extraction and Antimicrobial Properties of Azadirachta indica (Neem). Global Journal of Pharmacology. 2013; 7: 316-320.
- [12] Webb D. Phytochemicals' role in good health. Today's Dietitian. 2013; 15: 70.
- [13] Zillich OV, Schweiggert-Weisz U, Eisner P, Kerscher M. Polyphenols as active ingredients for cosmetic products. International Journal of Cosmetic Science. 2015; 37: 455-464.
- [14] Akagbue BO, Ibrahim MN, Ofure OF, Ekugbe OU, Amaobichukwu CT, et al. Comprehensive Assessment and Remediation Strategies for Air Pollution: Current Trends and Future Prospects; A Case Study in Bompai Industrial Area, Kano State, Nigeria. Communication in Physical Sciences. 2023a; 10: 1-13.
- [15] Abdulbariu I, Omeneke AL, Baba Aminu M, Dung PD, Salisu SM, et al. Application of Vertical Electrical Sounding (VES) for the Determination of Water-Bearing Zone in Karaworo, Lokoja Kogi State, Nigeria. Journal of Geography, Environment and Earth Science International. 2023; 27: 47-73.
- [16] Akagbue BO, Baba Aminu M. Effect of Na²², Cl-36, 3-H, and P-32 Exposure on Laboratory Clinical Researchers. Communication in Physical Sciences. 2023b; 9: 438- 446.
- [17] Biswas K, Chattopadhyay I, Banerjee RK, Bandyopadhyay U. Biological activities and medicinal properties of neem

Citation: Tasha Siame, Olayiwola Bolaji, Yisa Adeniyi Abolade, Sarah Aduwa, Amarachukwu Bernaldine, et al. (2025) Phytochemical Screening of *Azadirachta Indica* (Neem) Leaves: Revealing Health and Environmental Benefits. *Applied Medical Research*. AMR-1047

- (*Azadirachta indica*). *Current Science*. 2012; 82: 1336-1345.
- [18] Hossain MA, Shah MD, Sakari M. "Gas chromatography–mass spectrometry analysis of various organic extracts of *Merremia borneensis* from Sabah," *Asian Pacific Journal of Tropical Medicine*. 2011; 4: 637-641.
- [19] Ramesh SS, Suganya T. A review on phytochemical, pharmacological properties of neem plant. *World Journal of Pharmacy and Pharmaceutical Sciences*. 2018; 7: 85-97.
- [20] Kikani BA, Kikani SB, Panara NK. Neem (*Azadirachta indica*): Its potential for prevention and treatment of cancer. *Recent Patents on Anti-Cancer Drug Discovery*. 2020; 15: 256- 269.
- [21] Verma AK, Kulkarni P, Varma A. Potential herbs used as antidandruff in Ayurveda. *World Journal of Pharmacy and Pharmaceutical Sciences*. 2017; 6: 1632-1641.
- [22] Gupta S, Shukla R. In vitro antioxidant effect of various parts of *Azadirachta indica*. *International Journal of Pharmaceutical Sciences and Research*. 2014; 5: 1344-1348.
- [23] Isman MB. *Azadirachtin* and neem oil as natural insecticides: History, production, and perspectives. *Integrated Control in Oilseed Crops*. 2021; 43-57.
- [24] Dwivedi C, Dubey MP. Triterpenoids from *Azadirachta indica*: Chemistry and biological activity. *Studies in Natural Products Chemistry*. 2007; 34: 205-248.
- [25] Cui L, Zhang L, Jiang R, Zhang Y, Li M, et al. A review: The bioactivities and pharmacological applications of beta-sitosterol. *PLoS ONE*. 2018; 13: e0192307.
- [26] Dasgupta BR, Rao MR. Medicinal importance of tannins: An overview. *International Journal of Biomedical Research*. 2018; 9: 561-564.
- [27] Subapriya R, Nagini S. Medicinal properties of neem leaves: a review. *Current Medicinal Chemistry*. 2015; 12: 1387-1395.
- [28] Pazyar N, Yaghoobi R, Bagherani N, Kazerouni A. A review of applications of tea tree oil in dermatology. *International Journal of Dermatology*. 2012; 52: 784-790.
- [29] Chatterjee A, Pakrashi S. *The Treatise on Indian Medicinal Plants*. National Institute of Science Communication. 1991.
- [30] Adeniyi A, Bello I, Mukaila T, Hammed A. A Review of Microbial Molecular Profiling during Biomass Valorization. *Biotechnol Bioproc*. 2022; 27: 515–532.
- [31] Kumar SS, Dubey M. Phytoremediation: An eco-sustainable green technology for pollution prevention, control, and remediation. In *Handbook of Research on Inventive Bioremediation Techniques*. 2014; 155-188.
- [32] Patil PR, Patil SP, Mane A, Verma S. "Antidiabetic activity of alcoholic extract of Neem (*Azadirachta indica*) root bark," *National Journal of Physiology, Pharmacy and Pharmacology*. 2013; 3: 142-146.
- [33] Dholi SK, Raparla RS, Mankala K, Nagappan K. "Invivo antidiabetic evaluation of Neem leaf extract in alloxan induced rats," *Journal of Applied Pharmaceutical Science*. 2011; 1: 100-105.
- [34] Jeyasankar A, Amirtham D. Phytochemical and pharmacological profile of *Azadirachta indica*. *Indian Journal of Natural Products and Resources*. 2014; 5: 267-273.
- [35] Baba Aminu M, Kabiru GI, Simon DC, Changde AN, Andarawus Y, et al. Interrogating the Effects of Sand Mining: A Case Study of Agila District, Ado Local Government Area, Benue State, Nigeria, *Fudma Journal of Sciences*. 2023a; 20237: 317-331.
- [36] Akagbue BO, Aminu MB, Siame T, Ofure OF, Amaobichukwu CT, et al. Medical Effect of Emission from Unregulated Refineries at Rumuolumeni Community in Port Harcourt Rivers State Nigeria, *Asian Journal of Medicine and Health*. 2023c; 21: 83-94.
- [37] Edeoga HO, Okwu DE, Mbaebie BO. Phytochemical constituents of some Nigerian medical plants, *African Journal of Biotechnology*. 2005; 4: 685-688.
- [38] Madaki FM, Kabiru AY, Bakare Odunola AY, Mailafiya MT, Hamzah SC, et al. Phytochemical and Proximate Analyses of Methanol Leaf Extract of Neem, *Azadirachta indica*. 2016; 7: 45-67
- [39] Emran TB, Nasir Uddin MM, Rahman A, Uddin Z, Islam M. Phytochemical, Antimicrobial, Cytotoxic, Analgesic and AntiInflammatory Properties of *Azadirachta Indica*: A Therapeutic Study, *Journal of Bioanalysis and Biomedicine*. 2015; 12.
- [40] Ahmad M, Amin S. Phytochemical analysis and comprehensive evaluation of cardioprotective, antioxidant, and antimicrobial potential of *Azadirachta indica* leaves, *BioMedical Research International*. 2014; 301983.
- [41] Sajid I, Zafar M, Yasmin A. Recent Advances in Alkaloid Metabolism: Alkaloid Diversity and Evolution, *Molecules*. 2021; 26: 360.
- [42] Oleszek W, Stochmal A, Janda B, Piacente S. Saponins in the genus *Medicago* L, *Phytochemistry*. 2011; 56: 537-545.
- [43] Kumar R, Sharma S, Devi L. Investigation of Total Phenolic, Flavonoid Contents and Antioxidant Activity from Extracts of *Azadirachta indica* of Bundelkhand Region, *International Journal of Life Sciences Scientific Research*. 2018; 4: 1925-1933.
- [44] Muhammad IA, Aliyu LS, Ismalia A, Musa A. Nematicidal Properties and Phytochemical Screening of *Tamarindus Indica*

- and *Azadirachta Indica* Extract Against Burrowing Nematode of Banana Plant in Sokoto State, *Journal of Advanced Botany and Zoology*. 2018; V5I3.
- [45] López R. Saponins from the Spanish Arid Zone: a Review of Sources, Composition and Bioactivities, *Molecules*. 2015; 20: 13159-13194.
- [46] Schönthal AH. Direct Effects of Saponins on Cancer Cell Viability. In *Natural Compounds as Inducers of Cell Death*, Springer. 2018; 149-164.
- [47] Liu Y, Xu F, Jin C, Wu S. Saponins from Chinese Medicines as Anticancer Agents, *Molecules*. 2016; 21: 1326.
- [48] Salehi B, Valussi M, Jugran AK, Martorell M, Ramírez Alarcón K, et al. Phytochemicals in *Helicobacter pylori* Infections: What Are We Doing Now?, *International Journal of Molecular Sciences*. 2019; 20: 1254.
- [49] D'Archivio M, Filesi C, Vari R, Scazzocchio B, Masella R. Bioavailability of the polyphenols: status and controversies, *International Journal of Molecular Sciences*. 2010; 8: 1153-1175.
- [50] Visioli F, Caruso D, Galli C, Viappiani S. Polyphenols and human health: a prospectus, *Critical Reviews in Food Science and Nutrition*. 2011; 51: 524-546.
- [51] Gu L, Kelm MA, Hammerstone JF, Beecher G, Holden J, et al. Concentrations of proanthocyanidins in common foods and estimations of normal consumption, *Journal of Nutrition*. 2017; 134: 613-617.
- [52] Chen Z, Zhang P, Che Y, Wang L. Cardiotoxic steroids: Novel cancer therapeutic targets, *Oncotarget*. 2015; 6: 18206-18220.
- [53] Wu Y, Xu J, Xu J, Zheng W, Chen Q, et al. Cardiac glycosides stimulate Ca²⁺ increases and apoptosis in androgen-independent, metastatic human prostate adenocarcinoma cells, *Cancers*. 2020; 12: 220.
- [54] Zhang Y, Li J, Ji Y, Zhao T, Yang W. Cardiac glycosides suppress the maintenance of stemness and malignancy via inhibiting HIF-1 α in breast cancer stem cells, *Anti-Cancer Agents in Medicinal Chemistry*. 2020; 20: 71-80.
- [55] Grossman Y, Yu J, Byron KL, Bystroff C. In silico study of toxic cardiac glycoside binding to the Na⁺/K⁺ pump, *Biophysical Journal*. 2017; 113: 1063-1071.
- [56] Chan L, Anderson S, Patel R, Gerkin RD, Yoon SH. Reversal of severe digoxin overdose with digoxin-specific antibody, *The American Journal of Emergency Medicine*. 2020; 38: 1712-1713.
- [57] Yeh CK, Huang SM, Hwang KL, Huang YS. Acute ingestion of oleander leaf extracts induces apoptosis in HeLa cells, *Toxins*. 2019; 11: 16.
- [58] Scalbert A. Antimicrobial properties of tannins, *Phytochemistry*. 1991; 30: 3875-3883.
- [59] Cushnie TP, Lamb AJ. Antimicrobial activity of flavonoids, *International Journal of Antimicrobial Agents*. 2015; 26: 343-356.
- [60] Kasote DM, Hegde MV, Khedkar CD, Bae H. Significance of antioxidant potential of plants and its relevance to therapeutic applications, *International Journal of Biological Sciences*. 2015; 11: 982-991.
- [61] He L, Mu C, Shi J, Zhang Q, Shi B, et al. Comparison of tea tannin and teeth staining between green tea and black tea in vitro, *Journal of Dentistry*. 2018; 74: 35-39.
- [62] Rice Evans CA, Miller NJ, Paganga G. Antioxidant properties of phenolic compounds, *Trends in Plant Science*. 2007; 2: 152-159.
- [63] Dash SP, Dixit S, Sahoo S. Phytochemical and Biochemical Characterizations from Leaf Extracts from *Azadirachta indica*: An Important Medicinal Plant, *Biochemistry and Analytical Biochemistry*. 2017; 6: 323.
- [64] Saiyam R. Evaluation of *Azadirachta indica* and *Butea frondosa* extracts against gastrointestinal nematodosis in goats, MV Sc thesis (Parasitology) Nanaji Deshmukh Veterinary Science University Jabalpur. 2018.
- [65] Boyer J, Liu RH. Apple phytochemicals and their health benefits, *Nutrition Journal*. 2014 3: 5.
- [66] Mertens Talcott SU, Talcott ST, Percival SS. Low concentrations of quercetin and ellagic acid synergistically influence proliferation, cytotoxicity, and apoptosis in MOLT-4 human leukemia cells, *The Journal of Nutrition*. 2018; 133: 2669-2674.
- [67] Nishino H, Murakoshi M, Ii T, Takemura M, Kuchide M, et al. Carotenoids in cancer chemoprevention, *Cancer and Metastasis Reviews*. 2019; 19: 259-265.
- [68] Raut JS, Lakkakula S. Anti-inflammatory and antimicrobial activities of terpenes from Aromatic and Medicinal Plants, *Journal of Essential Oil-Bearing Plants*. 2019; 22: 1535- 1547.
- [69] Kim HM, Oh HA, Kim JH. Aromatherapy massage affects menopausal symptoms in Korean climacteric women: a pilot-controlled clinical trial, *Evidence Based Complementary and Alternative Medicine*. 2016; 23-65.
- [70] Foti C, Nettis E, Panebianco R, Calogiuri GF, Di Leo E, et al. Contact sensitization to fragrances in consecutive referred patients: a retrospective study, *Contact Dermatitis*. 2015; 49: 321-325.

Citation: Tasha Siame, Olayiwola Bolaji, Yisa Adeniyi Abolade, Sarah Aduwa, Amarachukwu Bernaldine, et al. (2025) Phytochemical Screening of Azadirachta Indica (Neem) Leaves: Revealing Health and Environmental Benefits. Applied Medical Research. AMR-1047

- [71] Musani SK, Mazi Kotwal N, Miller WL, Schoneveld O. Cardiovascular risk reduction, US Patent No. 2018; US20110171238A1.
- [72] Bouic PJ. The role of phytosterols and phytosterolins in immune modulation: a review of the past 10 years, Current Opinion in Clinical Nutrition and Metabolic Care. 2015; 4: 471-475.
- [73] Ostlund RE. Phytosterols in human nutrition, Annual Review of Nutrition. 2014; 22: 533- 549.
- [74] Pascual M, de Lorenzo C, García R, Suárez J, Cisteró A. Sterol content and isomer composition of phytosterol-enriched foods, Journal of Agricultural and Food Chemistry. 2016; 56: 3188-3192.
- [75] Alzohairy MA. Therapeutics role of Azadirachta indica (Neem) and their active constituents in diseases prevention and treatment, Evidence-Based Complementary and Alternative Medicine. 2016.
- [76] Maria C, Romilly MM. Phytochemical screening and the antimicrobial activity of the leaves of Azadirachta indica, International Journal of Scientific and Engineering Research. 2017; 8: 721-724.